




BIOCHEMISTRY REVIEW


Overview of Biomolecules

Chapter 13

Protein Synthesis






ABC
DEF
GHI
JKL
Base sequence
aa1
aa2
aa3
aa4
Amino acid sequence




Nonoverlapping code
A U A C G A G U C
1
2
3
Overlapping code
A U A C G A G U C
1
2
3

TABLE 27–1 Incorporation of Amino Acids into Polypeptides in Response to Random Polymers of RNA

Amino acid	*Observed frequency of incorporation (Lys = 100)*	*Tentative assignment for nucleotide composition* of corresponding codon*	*Expected frequency of incorporation based on assignment (Lys = 100)*
Asparagine	24	A_2C	20
Glutamine	24	A_2C	20
Histidine	6	AC_2	4
Lysine	100	AAA	100
Proline	7	AC_2, CCC	4.8
Threonine	26	A_2C, AC_2	24

Note: Presented here is a summary of data from one of the early experiments designed to elucidate the genetic code. A synthetic RNA containing only A and C residues in a 5:1 ratio directed polypeptide synthesis, and both the identity and the quantity of incorporated amino acids were determined. Based on the relative abundance of A and C residues in the synthetic RNA, and assigning the codon AAA (the most likely codon) a frequency of 100, there should be three different codons of composition A_2C, each at a relative frequency of 20; three of composition AC_2, each at a relative frequency of 4.0; and CCC at a relative frequency of 0.8. The CCC assignment was based on information derived from prior studies with poly(C). Where two tentative codon assignments are made, both are proposed to code for the same amino acid.

*These designations of nucleotide composition contain no information on nucleotide sequence (except, of course, AAA and CCC).


First letter of codon (5′ end)
Second letter of codon
U
C
A
G
UUU Phe
UUC Phe
UUA Leu
UUG Leu
UCU Ser
UCC Ser
UCA Ser
UCG Ser
UAU Tyr
UAC Tyr
UAA Stop
UAG Stop
UGU Cys
UGC Cys
UGA Stop
UGG Trp
CUU Leu
CUC Leu
CUA Leu
CUG Leu
CCU Pro
CCC Pro
CCA Pro
CCG Pro
CAU His
CAC His
CAA Gln
CAG Gln
CGU Arg
CGC Arg
CGA Arg
CGG Arg
AUU Ile
AUC Ile
AUA Ile
AUG Met
ACU Thr
ACC Thr
ACA Thr
ACG Thr
AAU Asn
AAC Asn
AAA Lys
AAG Lys
AGU Ser
AGC Ser
AGA Arg
AGG Arg
GUU Val
GUC Val
GUA Val
GUG Val
GCU Ala
GCC Ala
GCA Ala
GCG Ala
GAU Asp
GAC Asp
GAA Glu
GAG Glu
GGU Gly
GGC Gly
GGA Gly
GGG Gly



Reading frame 1
5′---U U C U C G G A C C U G G A G A U U C A C A G U---3′
Reading frame 2
---U U C U C G G A C C U G G A G A U U C A C A G U---
Reading frame 3
---U U C U C G G A C C U G G A G A U U C A C A G U---

TABLE 27–3 Degeneracy of the Genetic Code

Amino acid	*Number of codons*	*Amino acid*	*Number of codons*
Met	1	Tyr	2
Trp	1	Ile	3
Asn	2	Ala	4
Asp	2	Gly	4
Cys	2	Pro	4
Gln	2	Thr	4
Glu	2	Val	4
His	2	Arg	6
Lys	2	Leu	6
Phe	2	Ser	6

TABLE 5.5 **Distinctive codons of human mitochondria**

Codon	Standard code	Mitochondrial code
UGA	Stop	Trp
UGG	Trp	Trp
AUA	Ile	Met
AUG	Met	Met
AGA	Arg	Stop
AGG	Arg	Stop

Wild type allele:

M D D Q S R M L Q T L A G V N L

atggacgatcaatccaggatgctgcagactctggccggggtgaacctg

silent (third base pair) mutation:

M D D Q S R M L Q T L A G V N L

atggacgatcaatccaggatgctgcaaactctggccggggtgaacctg

point mutation (missense):

M D D Q S R M L K T L A G V N L

atggacgatcaatccaggatgctgaagactctggccggggtgaacctg

point mutation (nonsense):


M D D Q S R M L stop

atggacgatcaatccaggatgctgtagactctggccggggtgaacctg

frameshift leading to premature termination:

M D D Q S R M L R L W P G stop

atggacgatcaatccaggatgctgagactctggccggggtgaacctg


mRNA
5′
G U A G C C U A C G G A U
3′
(+)
Insertion
G U A G C C U C A C G G A U
(–)
Deletion
G U A C C U A C G G A U
(+)
(–)
Insertion and deletion
G U A A G C C A C G G A U
Reading frame restored



Are You Getting It??



Which properties are characteristic of the normal genetic code? ***(multiple answers)***

a) A nucleotide in an mRNA molecule can be part of only one codon.
b) A codon can code for only one amino acid.
c) Each amino acid has only one codon.
d) There is one specific start codon.
e) There is one specific stop codon.
f) Every change in a codon produces a change in the amino acid sequence of the protein.



Are You Getting It??



Answer

Which properties are characteristic of the normal genetic code?

a) ***A nucleotide in an mRNA molecule can be part of only one codon.***

b) ***A codon can code for only one amino acid.***

c) **Each amino acid has only one codon.**

d) ***There is one specific start codon.***

e) **There is one specific stop codon.**


f) **Every change in a codon produces a change in the amino acid sequence of the protein.**


TABLE 5.2 **RNA molecules in *E. coli***


Type	Relative amount (%)	Sedimentation coefficient (S)	Mass (kd)	Number of nucleotides
Ribosomal RNA (rRNA)	80	23	1.2×10^3	3700
		16	0.55×10^3	1700
		5	3.6×10^1	120
Transfer RNA (tRNA)	15	4	2.5×10^1	75
Messenger RNA (mRNA)	5		Heterogeneous	


TABLE 27–5 Components Required for the Five Major Stages of Protein Synthesis in *E. coli*


Stage	*Essential components*
1. Activation of amino acids	20 amino acids
	20 aminoacyl-tRNA synthetases
	32 or more tRNAs
	ATP
	Mg^{2+}
2. Initiation	mRNA
	N-Formylmethionyl-$tRNA^{fmet}$
	Initiation codon in mRNA (AUG)
	30S ribosomal subunit
	50S ribosomal subunit
	Initiation factors (IF-1, IF-2, IF-3)
	GTP
	Mg^{2+}
3. Elongation	Functional 70S ribosome (initiation complex)
	Aminoacyl-tRNAs specified by codons
	Elongation factors (EF-Tu, EF-Ts, EF-G)
	GTP
	Mg^{2+}
4. Termination and release	Termination codon in mRNA
	Release factors (RF-1, RF-2, RF-3)
5. Folding and posttranslational processing	Specific enzymes, cofactors, and other components for removal of initiating residues and signal sequences, additional proteolytic processing, modification of terminal residues, and attachment of phosphate, methyl, carboxyl, carbohydrate, or prosthetic groups



amino acid
3′
A
C
C
5′
X
Y
Z
Anticodon


3′
5′
tRNA
Anticodon
3 2 1
U A G
mRNA 5′
A U C
3′
1 2 3
Codon


3′
Site for amino acid attachment
5′
pG
m1G
m2G
m1I
I
5′
3′
Anticodon triplet


3′
A
C
C
Pu
Amino acid arm
5′
pG
D arm
Pu
A
G*
G
A
U
TψC arm
C
G
T
ψ
C
Contains two or three D residues at different positions
Extra arm
Variable in size, not present in all tRNAs
Py
U
Pu
Anticodon arm
Wobble position
5′
3′
Anticodon


Amino
acid arm
TψC arm
64
54
1
5′
56
72
3′
D arm
(residues
10–25)
7
69
20
12
44
26
Anticodon
arm
38
Anticodon
32
(a)


(b)



Are You Getting It??



Which properties are found in tRNA molecules?
(multiple answers)

a) They contain 1000-2000 nucleotides.
b) They are single-stranded nucleic acids.
c) They have secondary and tertiary structure.
d) The bases are always A, C, G, and U.
e) They contain hydrogen bonds.
f) They contain three specific bases that form a codon.
g) They can carry an amino acid at the 3'- end.



Are You Getting It??



Answer


Which properties are found in tRNA molecules?


a) **They contain 1000-2000 nucleotides.**
b) ***They are single-stranded nucleic acids.***
c) ***They have secondary and tertiary structure.***
d) **The bases are always A, C, G, and U.**
e) ***They contain hydrogen bonds.***
f) **They contain three specific bases that form a codon.**
g) ***They can carry an amino acid at the 3'- end.***


OVERALL REACTION


amino acid + tRNA + ATP


aminoacyl-tRNA + AMP + PP_i


3′ end of tRNA
Adenine
H
2′
OH
H
O
H
3′
O
C
C
R
H
CH2
H
O
O
Aminoacyl group
−O—P=O
+NH3
O
5′
pG
Amino acid arm
D arm
TψC arm
Anticodon arm



Amino acid
ATP
Adenine
1
PPi
Adenine
5'-Aminoacyl adenylate
(aminoacyl-AMP)


Adenine
5'-Aminoacyl adenylate
(aminoacyl-AMP)
class I
aminoacyl-tRNA
synthetases
class II
aminoacyl-tRNA
synthetases
3' end of tRNA
Adenosine
Aminoacyl-AMP
tRNA
Adenosine
Aminoacyl-AMP
tRNA



3' end of tRNA
Adenine
Adenosine
Aminoacyl-AMP
tRNA
2a
AMP
2b
AMP
transesterification
3a
Aminoacyl-tRNA




Valine



Isoleucine


Editing site
Activation site


3′
Amino acid arm
5′
TψC arm
D arm
Extra arm
Anticodon arm
Anticodon



Are You Getting It??



Which mechanisms are characteristic of aminoacyl-tRNA synthetases? ***(multiple answers)***

a) They form a covalent bond between an amino acid and a tRNA molecule.
b) They break down ATP for energy.
c) One enzyme can react with all twenty amino acids.
d) They have two active sites to proof-read the amino acid.
e) They recognize specific nucleotides in tRNA molecules.



Are You Getting It??



Answer

Which mechanisms are characteristic of aminoacyl-tRNA synthetases?


a) ***They form a covalent bond between an amino acid and a tRNA molecule.***

b) ***They break down ATP for energy.***

c) **One enzyme can react with all twenty amino acids.**

d) ***They have two active sites to proof-read the amino acid.***

e) ***They recognize specific nucleotides in tRNA molecules.***


3′
5′
tRNA
Anticodon
3 2 1
U A G
mRNA 5′
A U C
3′
1 2 3
Codon



3 2 1
3 2 1
3 2 1
Anticodon (3′) G–C–I G–C–I G–C–I (5′)
Codon (5′) C–G–A C–G–U C–G–C (3′)
1 2 3
1 2 3
1 2 3

TABLE 27–4 How the Wobble Base of the Anticodon Determines the Number of Codons a tRNA Can Recognize

1. One codon recognized:

Anticodon	(3′) X–Y–**C** (5′)	(3′) X–Y–**A** (5′)
Codon	(5′) Y–X–**G** (3′)	(5′) Y–X–**U** (3′)

2. Two codons recognized:

Anticodon	(3′) X–Y–**U** (5′)	(3′) X–Y–**G** (5′)
Codon	(5′) Y–X–**A** or **G** (3′)	(5′) Y–X–**C** or **U** (3′)

3. Three codons recognized:

Anticodon	(3′) X–Y–**I** (5′)
Codon	(5′) Y–X–**A** or **U** or **C** (3′)

Note: X and Y denote bases complementary to and capable of strong Watson-Crick base pairing with X′ and Y′, respectively. Wobble bases—in the 3′ position of codons and 5′ position of anticodons—are shaded in pink.

TABLE 29.3 Allowed pairings at the third base of the codon according to the wobble hypothesis

First base of anticodon	Third base of codon
C	G
A	U
U	A or G
G	U or C
I	U, C, or A



Are You Getting It??



Which kinds of interactions can occur between a codon and an anticodon? ***(multiple answers)***

a) They bind to each other non-covalently.
b) The three base- pairs formed have equal strength.
c) They bind in a antiparallel orientation.
d) The first base in a codon can wobble.
e) The first base in an anticodon can be a rare base.
f) An anticodon can bind to only one codon.



Are You Getting It??



Answer

Which kinds of interactions can occur between a codon and an anticodon?

a) ***They bind to each other non-covalently.***


b) **The three base- pairs formed have equal strength.**


c) ***They bind in a antiparallel orientation.***

d) **The first base in a codon can wobble.**

e) ***The first base in an anticodon can be a rare base.***

f) **An anticodon can bind to only one codon.**


(A)
50 nm
(500 Å)
(B)
(C)


Bacterial ribosome
70S
M_r 2.7 × 10⁶
50S
M_r 1.8 × 10⁶
5S rRNA
(120 nucleotides)
23S rRNA
(3,200 nucleotides)
36 proteins
30S
M_r 0.9 × 10⁶
16S rRNA
(1,540 nucleotides)
21 proteins
Eukaryotic ribosome
80S
M_r 4.2 × 10⁶
60S
M_r 2.8 × 10⁶
5S rRNA
(120 nucleotides)
28S rRNA
(4,700 nucleotides)
5.8S rRNA
(160 nucleotides)
~ 49 proteins
40S
M_r 1.4 × 10⁶
18S rRNA
(1,900 nucleotides)
~ 33 proteins




50S subunit
70S ribosome
30S subunit

TABLE 27-6 RNA and Protein Components of the *E. coli* Ribosome

Subunit	*Number of different proteins*	*Total number of proteins*	*Protein designations*	*Number and type of rRNAs*
30S	21	21	S1-S21	1 (16S rRNA)
50S	33	36	L1-L36*	2 (5S and 23S rRNAs)

*The L1 to L36 protein designations do not correspond to 36 different proteins. The protein originally designated L7 is in fact a modified form of L12, and L8 is a complex of three other proteins. Also, L26 proved to be the same protein as S20 (and not part of the 50S subunit). This gives 33 different proteins in the large subunit. There are four copies of the L7/L12 protein, with the three extra copies bringing the total protein count to 36.


5′(1)
3′ (1,542)
16S rRNA
5′
3′
5S rRNA



Are You Getting It??



Which component is found in E. coli ribosomes?

a) They contain two subunits of equal size.
b) They contain three types of rRNAs.
c) They contain over 100 different proteins.
d) They contain codons.
e) They contain anticodons.
f) They contain aminoacyl-tRNA synthetases.




Are You Getting It??



Answer

Which component is found in E. coli ribosomes?

a) They contain two subunits of equal size.
b) *They contain three types of rRNAs.*
c) They contain over 100 different proteins.
d) They contain codons.
e) They contain anticodons.
f) They contain aminoacyl-tRNA synthetases.


5'
TCA
GAG
GTG
AAA
TGC
TAT
GGT
3'
DNA coding strand
3'
AGT
CTC
CAC
TTT
ACG
ATA
CCA
5'
DNA Template strand
Transcription
5'
UCA
GAG
GUG
AAA
UGC
UAU
GGU
3'
mRNA
Translation
N
Ser
Glu
Val
Lys
Cys
Tyr
Gly
C
Peptide chain

INITIATION COMPLEX

mRNA


30 S subunit

50 S subunit


GTP


Initiation Factors


First tRNA




***N*-Formylmethionine**



tRNAf
+
Methionine
Synthetase
H2N
S
CH3
H
O
O
tRNAf
Methionyl-tRNAf
(Met-tRNAf)
N10-Formyl-
tetrahydrofolate
Transformylase
Tetrahydrofolate
O
NH
H
S
CH3
H
O
O
tRNAf
Formylmethionyl-tRNAf
(fMet-tRNAf)



IF-3
IF-1
30S
Subunit
P
A
mRNA
1
Initiation
codon
mRNA
5′
A U G
3′
IF-3
P
IF-1



(A) **Prokaryotic start signal**


E. coli trpA (5′) A G C A C G A G G G G A A A U C U G A U G G A A C G C U A C (3′)
E. coli araB U U U G G A U G G A G U G A A A C G A U G G C G A U U G C A
E. coli lacI C A A U U C A G G G U G G U G A A U G U G A A A C C A G U A
ϕX174 phage A protein A A U C U U G G A G G C U U U U U U A U G G U U C G U U C U
λ phage cro A U G U A C U A A G G A G G U U G U A U G G A A C A A C G C
Shine-Dalgarno sequence; pairs with 16S rRNA
Initiation codon; pairs with fMet-tRNA fMet
(a)
Prokaryotic mRNA with consensus Shine-Dalgarno sequence
3′ End of 16S rRNA
3′OH
A U U C C U C C A
G A U C A
(5′) G A U U C C U A G G A G G U U U G A C C U A U G C G A G C U U U U A G U (3′)
(b)


1
Initiation codon
mRNA
5′
A U G
3′
IF-3
P
IF-1
2
5′
fMet
IF-2
GTP
U A C
(3′) UAC (5′)
Anticodon
5′
fMet
tRNA
IF-2
GTP
U A C
5′
A U G
3′
IF-3
P
IF-1


5′
fMet
U A C
(3′) UAC (5′)
Anticodon
5′
fMet
tRNA
IF-2 – GTP
U A C
5′
A U G
3′
IF-3
P
IF-1
50S
Subunit
GDP + P_i
IF-1 + IF-2 + IF-3
③
5′
fMet
50S
Subunit
E
U A C
5′
A U G
3′
P
A
Next
codon



Are You Getting It??



Which events occur during the initiation of translation in E. coli? ***(multiple answers)***

a) **The first codon recognized is always UAG.**
b) **The first amino acid used is always N-formylmethionine.**
c) **The initiation factors needed are RNA molecules.**
d) **The first tRNA molecule binds to the P site.**
e) **The 16S rRNA correctly positions the mRNA.**
f) **GTP is broken down as the ribosome forms.**



Are You Getting It??



Answer

Which events occur during the initiation of translation in E. coli?

a) **The first codon recognized is always UAG.**
b) ***The first amino acid used is always N-formylmethionine.***
c) **The initiation factors needed are RNA molecules.**
d) ***The first tRNA molecule binds to the P site.***
e) ***The 16S rRNA correctly positions the mRNA.***
f) ***GTP is broken down as the ribosome forms.***





40S Ribosomal subunit
5′cap
eIF3
3′poly(A) tail
A
A
A
A(A)n
eIF4E
eIF4G
PAB
AUG
Gene
3′Untranslated region


TABLE 27–8 Protein Factors Required for Initiation of Translation in Bacterial and Eukaryotic Cells


Factor	*Function*
Bacterial	
IF-1	Prevents premature binding of tRNAs to A site
IF-2	Facilitates binding of fMet-tRNA^{fMet} to 30S ribosomal subunit
IF-3	Binds to 30S subunit; prevents premature association of 50S subunit; enhances specificity of P site for fMet-tRNA^{fMet}
Eukaryotic*	
eIF2	Facilitates binding of initiating Met-tRNA^{Met} to 40S ribosomal subunit
eIF2B, eIF3	First factors to bind 40S subunit; facilitate subsequent steps
eIF4A	RNA helicase activity removes secondary structure in the mRNA to permit binding to 40S subunit; part of the eIF4F complex
eIF4B	Binds to mRNA; facilitates scanning of mRNA to locate the first AUG
eIF4E	Binds to the 5′ cap of mRNA; part of the eIF4F complex
eIF4G	Binds to eIF4E and to poly(A) binding protein (PAB); part of the eIF4F complex
eIF5	Promotes dissociation of several other initiation factors from 40S subunit as a prelude to association of 60S subunit to form 80S initiation complex
eIF6	Facilitates dissociation of inactive 80S ribosome into 40S and 60S subunits


*The prefix "e" identifies these as eukaryotic factors.



50S
Tunnel
E
A
P
E
P
A
30S
Aminoacyl-tRNA
binding
Peptide-bond
formation
GTP
Translocation
Elongation
factor G
GDP + Pi
tRNA
dissociation



Initiation complex
50S
5′
fMet
E
U A C
A U G
5′
3′
P
A
Initiation codon
Next codon
30S
Incoming aminoacyl-tRNA
5′
AA2
Tu
GTP
Ts
GDP
Pi
binding of incoming aminoacyl-tRNA


E site
P site
A site
fMet-tRNA fMet
Aminoacyl-
tRNA2
H
C=O
NH
R1–C–H
O=C
O
NH2
R2–C–H
O=C
O
5′
5′
E
U A C
A U G
mRNA 5′
3′
P
A


E site
P site
A site
Deacylated tRNA fMet
Dipeptidyl-tRNA2
OH
5′
5′
5′
5′
E
U A C
A U G
5′
3′
P
A


translocation
EF-G – GTP
EF-G + GDP + P_i
E site
P site
A site
Incoming aminoacyl-tRNA$_3$
5′
3′
U A C
A U G
P
A
Direction of ribosome movement


Release factor
binds
5′
RF
E
5′
U A G
3′
P
A
polypeptidyl-tRNA
link hydrolyzed
COO⁻
5′
RF
E
5′
U A G
3′
P
A


COO⁻
5′
RF
E
5′
U A G
3′
P
A
components
dissociate
5′
RF
5′
U A G
3′



Are You Getting It??



Which reactions occur during translation in E. coli?
(multiple answers)

a) EF-Tu binds to an aminoacyl-tRNA.
b) The incoming tRNA binds to the A site.
c) A non-covalent bond forms between two amino acids.
d) The 23S rRNA catalyzes movement of the mRNA.
e) EF- Ts causes translocation to occur.
f) A protein releasing factor binds to a stop codon.



Are You Getting It??



Answer

Which reactions occur during translation in E. coli?

a) ***EF-Tu binds to an aminoacyl-tRNA.***


b) ***The incoming tRNA binds to the A site.***


c) **A non-covalent bond forms between two amino acids.**


d) **The 23S rRNA catalyzes movement of the mRNA.**


e) **EF- Ts causes translocation to occur.**

f) ***A protein releasing factor binds to a stop codon.***


Incoming ribosomal subunits
60S
40S
5′
+
NH3
Growing
polypeptide
+
NH3
Direction
of
Translation
+
NH3
+
NH3
3′
+
NH3



0.25 μm



DNA
duplex
RNA
polymerase
5′
3′
3′
5′
Ribosome
5′
mRNA
+NH3
+NH3
Direction of transcription
Direction of translation



Ribosomes
on mRNA
DNA


POST-TRANSLATIONAL MODIFICATIONS


- **PROTEIN FOLDING**
- **N- and C-TERMINAL REMOVAL OF AAs**
- **MODIFICATION OF AMINO ACIDS**
- **ADDITION OF PROSTHETIC GROUPS**
- **CLEAVAGE OF POLYPEPTIDE CHAIN**
- **DISULFIDE BONDS**
- **PROTEIN TARGETING**


Phosphoserine
Phosphothreonine
Phosphotyrosine
(a)


γ-Carboxyglutamate
(b)


Methyllysine
Dimethyllysine
Trimethyllysine
Methylglutamate
(c)


Signal sequence
2
1
5′ cap
mRNA
GUA
8
Ribosome cycle
SRP cycle
Cytosol
SRP
3
5
GDP + Pi
AAA(A)n
Ribosome receptor
Endoplasmic reticulum
GTP
4
6
7
Peptide translocation complex
SRP receptor
Signal peptidase
ER lumen


Nucleus
Rough ER
Smooth ER
Transport vesicle
Cis side
Sorting
Trans side
Golgi complex
Lysosome
Secretory Granule
Transport vesicles
Cytosol
Plasma membrane



Are You Getting It??



Which events can occur during or after translation in E. coli?
(multiple answers)

a) Multiple ribosomes can bind to one mRNA.
b) Translation can begin only after transcription is completed.
c) An mRNA molecule is translated only once.
d) The new protein must fold into its proper conformation.
e) Amino acids in the new protein can be modified.
f) Every new protein molecule binds to a prosthetic group.
g) All new proteins function in the cell cytoplasm.



Are You Getting It??



Answer

Which events can occur during or after translation in E. coli?

a) ***Multiple ribosomes can bind to one mRNA.***

b) **Translation can begin only after transcription is completed.**

c) **An mRNA molecule is translated only once.**

d) ***The new protein must fold into its proper conformation.***

e) ***Amino acids in the new protein can be modified.***

f) **Every new protein molecule binds to a prosthetic group.**

g) **All new proteins function in the cell cytoplasm.**